

Mouse Anti-CD11b [MD131]: MC0500, MC0500RTU7

Intended Use: For Research Use Only

Description: CD11b, also known as ITAM, Integrin alpha-M or MAC-1 alpha subunit or CR3 alpha chain belongs to the integrin alpha chain family; it is predominately presented in human myeloid cells, NK1 cells, monocytes, granulocytes and follicular dendritic cells. The alpha subunit of ITAM/beta-2 complex (CD11b/CD18, Mac-1), is a receptor for fibrinogen, factor X, and ICAM1. ITAM/beta-2 is implicated in adhesive interactions of monocytes, macrophages, and granulocytes. CD11b has been used as a common myeloid marker. CD11b is expressed in about 50% of acute myeloid leukemia (AML). In combination with CD117, CD11b is helpful in differentiating acute promyelocytic leukemia (CD11b negative) from recovering benign myeloid proliferation (CD11b positive, CD117 negative). In acute promyelocytic leukemia patients treated with all-trans retinoic acid or Arsenic trioxide (As₂O₃), CD11b is a marker for differentiating the induction of leukemia cells. CD11b is also expressed on microglia cells and involved in the development of neurodegenerative diseases.

Specifications:


Clone: MD131
 Source: Mouse
 Isotype: IgG2b/k
 Reactivity: Human
 Immunogen: Recombinant fragment aa941-1074 of human ITGAM protein
 Localization: Membrane
 Formulation: Purified antibody in PBS pH7.4, containing BSA and ≤ 0.09% sodium azide (NaN₃)
 Storage: Store at 2°- 8°C
 Applications: IHC
 Package:

Description	Catalog No.	Size
CD11b Concentrated	MC0500	1 ml
CD11b Prediluted	MC0500RTU7	7 ml

IHC Procedure*:

Positive Control Tissue: Spleen, leukemia
 Concentrated Dilution: 50-200
 Pretreatment: Tris EDTA pH9.0, 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human tonsil stained with-anti-CD11b using DAB

References:

1. Inhaled Submicron Particle Paclitaxel (NanoPac) Induces Tumor Regression and Immune Cell Infiltration in an Orthotopic Athymic Nude Rat Model of Non-Small Cell Lung Cancer. Verco J, et al. J Aerosol Med Pulm Drug Deliv 32:266-277, 2019.
2. Osteoclast-Like Cells in Aneurysmal Disease Exhibit an Enhanced Proteolytic Phenotype. Kelly MJ, et al. Int J Mol Sci 20:N/A, 2019.

Doc. 100-MC0500
Rev. A