

Mouse Anti-CD33 [PWS44]: MC0515, MC0515RTU7

Intended Use: For Research Use Only

Description: CD33 (gp67, or siglec-3) is a 67 kDa glycosylated transmembrane protein that is a member of the sialic acid-binding immunoglobulin-like lectin (siglec) family. This CD33 antibody may be particularly advantageous for cases of acute myeloid leukemia, minimally differentiated (AML-M0) and acute monocytic leukemia (AML-M5), in which other paraffin section markers of myeloid differentiation (such as anti-myeloperoxidase) may be negative. All cases of myeloid sarcoma in this study showed anti-CD33 positivity in the myeloid and monocytic subsets, allowing for easy interpretation. The excellent sensitivity and specificity for myelomonocytic lineage makes this anti-CD33 a useful diagnostic marker for myeloid sarcoma. In addition, this anti-CD33 may be useful in determining CD33 expression on previous paraffin-embedded material if flow cytometry studies were not initially performed in patients with acute leukemia. Analysis of CD33 expression in paraffin-embedded bone marrow biopsy specimens provides another alternative when evaluating acute leukemias.


Specifications

Description		Catalog No.	Size
Package:			
Applications:	IHC		
Storage:	Store at 2°- 8°C		
Formulation:	Antibody in PBS pH7.4, con	taining BSA and $\leq 0.09\%$ so	odium azide (NaN3)
Localization:	Membrane, cytoplasm		
Reactivity:	Human		
Isotype:	IgG2b		
Source:	Mouse		
Clone:	PWS44		

Description	Catalog No.	Size
CD33 Concentrated	MC0515	1 ml
CD33 Prediluted	MC0515RTU7	7 ml

IHC Procedure*

IIIC I I OCCUUI C			
Positive Control Tissue:	Tonsil, myeloid leukemia		
Concentrated Dilution:	10-50		
Pretreatment:	Citrate pH6.0 or EDTA pH8.0, 15 min Pressure Cooker or 30-60 min water bath at 95°-99°C		
Incubation Time and Temp:	30-60 minutes @ RT		
Detection:	Refer to the detection system manual		
* Result should be confirmed by an established diagnostic procedure.			



FFPE human Acute myeloid leukemia stained with anti-CD33 using DAB

References:

- 1. CD33-specific chimeric antigen receptor T cells exhibit potent preclinical activity against human acute myeloid leukemia. Kenderian SS, et al. Leukemia. Aug;29(8):1637-47, 2015.
- 2. Association of CD33 polymorphism rs3865444 with Alzheimer's disease pathology and CD33 expression in human cerebral cortex. Walker DG, et al. Neurobiol Aging. Feb;36(2):571-82, 2015.
- 3. Indolent T-lymphoblastic proliferation with disseminated multinodal involvement and partial CD33 expression. Ohgami RS, et al. Am J Surg Pathol. Sep;38(9):1298-304, 2014.

Doc. 100-MC0515 Rev. A