

**Rabbit Anti-LEF1/TCF1 Alpha [MD158R]: RM0195, RM0195RTU7**

**Intended Use:** For Research Use Only

**Description:** LEF1 or TCF1 alpha participates in the Wnt signaling pathway, activates transcription of target genes in the presence of CTNNB1 and EP300. It may play a role in hair cell differentiation and follicle morphogenesis. TLE1, TLE2, TLE3 and TLE4 repress transactivation mediated by LEF1 and CTNNB1. It regulates T-cell receptor alpha enhancer function. PIAG antagonizes both Wnt-dependent and Wnt-independent activation by LEF1. Isoform 3 lacks the CTNNB1 interaction domain and may be an antagonist for Wnt signaling. Isoform 5 transcriptionally activates the fibronectin promoter, binds to and represses transcription from the E-cadherin promoter in a CTNNB1-independent manner, and is involved in reducing cellular aggregation and increasing cell migration of pancreatic cancer cells. Isoform 1 transcriptionally activates MYC and CCND1 expression and enhances proliferation of pancreatic tumor cells. Detected in thymus but not detected in normal colon, but highly expressed in colon cancer biopsies and colon cancer cell lines. Expressed in several pancreatic tumors and weakly expressed in normal pancreatic tissue. Isoforms 1 and 5 are detected in several pancreatic cell lines.


**Specifications**

Clone: MD158R  
 Source: Rabbit  
 Isotype: IgG  
 Reactivity: Human  
 Immunogen: Recombinant fragment of human LEF1 protein around aa100-200  
 Localization: Nucleus  
 Formulation: Antibody in PBS pH7.4, containing BSA and ≤ 0.09% sodium azide (NaN3)  
 Storage: Store at 2°- 8°C  
 Applications: IHC  
 Package:

Description	Catalog No.	Size
LEF1/TCF1 Alpha Concentrated	RM0195	1 ml
LEF1/TCF1 Alpha Prediluted	RM0195RTU7	7 ml

**IHC Procedure**

Positive Control: Tonsil, colon carcinoma  
 Concentrated Dilution: 50-200  
 Pretreatment: Tris EDTA pH9.0, 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C  
 Incubation Time and Temp: 30-60 minutes @ RT  
 Detection: Refer to the detection system manual  
 \* Result should be confirmed by an established diagnostic procedure.



FFPE human lymph node stained with anti-LEF1 using DAB

**References:**

1. Expression of LEF1 is an independent prognostic factor for patients with oral squamous cell carcinoma. Su MC, et al. J Formos Med Assoc. Sep 7, 2013.
2. LEF-1 and TCF4 expression correlate inversely with survival in colorectal cancer. Kriegl L, et al. J Transl Med. Nov 22;8:123, 2010.