

Mouse Anti-Napsin A [MD46]: MC0133, MC0133RTU7

Intended Use: For Research Use Only

Description: Napsin A has a specific function in normal alveolar epithelium and is proposed to play a role in the proteolytic processing of surfactant precursors. Napsin A is reported to be predominantly expressed in lamellar bodies of type II pneumocytes, secondary lysosomes of alveolar macrophages, respiratory epithelium of terminal and respiratory bronchioles, plasma cells, within a subset of lymphocytes in normal lung, as well as in epithelial cells of renal tubules in normal kidney and is weakly expressed in normal spleen. Napsin A is an aspartic proteinase that belongs to the peptidase A1 family and plays a role in pneumocyte surfactant processing. In normal tissue, Anti-Napsin A specifically labels type II pneumocytes in adult lung and epithelial cells in kidney tissues. In abnormal tissues, Studies have reported that Napsin A is expressed in 90% of primary lung adenocarcinomas. Napsin A and 79% of renal cell carcinoma by immunohistochemistry. Napsin A is a useful marker for lung adenocarcinoma. The combined use of Napsin A and thyroid transcription factor (TTF) improves the sensitivity and specificity for identification of pulmonary adenocarcinoma.

Specifications


Clone:	MD46
Source:	Mouse
Isotype:	IgG1k
Reactivity:	Human
Immunogen:	Recombinant human Napsin A protein fragment aa189-299
Localization:	Cytoplasm
Formulation:	Purified antibody in PBS pH 7.4, containing BSA and ≤ 0.09% sodium azide (NaN3)
Storage:	Store at 2°- 8°C
Applications:	IHC
Package:	

Description	Catalog No.	Size
Napsin A Concentrated	MC0133	1 ml
Napsin A Prediluted	MC0133RTU7	7 ml

IHC Procedure*

Positive Control Tissue:	Lung carcinoma
Concentrated Dilution:	50-200
Pretreatment:	Tris EDTA pH9.0 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C
Incubation Time and Temp:	30-60 minutes @ RT
Detection:	Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human lung adenocarcinoma stained with anti-Napsin A using DAB

References:

1. Napsin A staining in adrenal cortical neoplasms. Ballard M, et al. Arch Pathol Lab Med. Jul;137(7):883, 2013.
2. Value of PAX8, PAX2, napsin A, carbonic anhydrase IX, and claudin-4 immunostaining in distinguishing pleural epithelioid mesothelioma from metastatic renal cell carcinoma. Ordóñez NG. Mod Pathol. Aug;26(8):1132-43, 2013.
3. Comparison of monoclonal napsin A, polyclonal napsin A, and TTF-1 for determining lung origin in metastatic adenocarcinomas. Mukhopadhyay S, et al. Am J Clin Pathol., Nov;138(5):703-11, 2012.