

Rabbit Anti-OCT4 [MD192R]: RM0148, RM0148RTU7

Intended Use: For Research Use Only

Description: OCT4, also known as OTF3 or POU5F1, is a member of the POU family of transcription factors, involved in the regulation of pluripotency during normal development and is detectable in embryonic stem and germ cells. It can specifically bind to the octamer motif (5'-ATTTTCAT-3'), and it is critical for the self-renewal of embryonic stem cells. Overall, OCT4 is a key regulator of self-renewal in embryonic stem cells; its expression is potentially correlated with tumorigenesis and can affect some aspects of tumor behavior such as tumor recurrence or resistance to therapies. OCT4 is expressed in undifferentiated pluripotency cells, germ cells in ovary and testes. OCT4 is a sensitive and specific marker for germ cell tumors. It is consistently detected in carcinoma in situ/gonadoblastoma, seminomas, germinoma, dysgerminoma, and embryonal carcinoma but not in the differentiated components of nonseminomas, i.e., teratomas, yolk sac tumors, and choriocarcinomas. An antibody to OCT4 is useful in the identification of primary as well as metastatic germ cell tumors.

Specifications


Clone: MD192R
 Source: Rabbit
 Isotype: IgG
 Reactivity: Human
 Immunogen: Recombinant fragment of human OCT-4/POU5F1 protein aa1-134
 Localization: Nucleus
 Formulation: Purified antibody in PBS pH7.4, containing BSA and ≤ 0.09% sodium azide (NaN₃)
 Storage: Store at 2°- 8°C
 Applications: IHC
 Package:

Description	Catalog No.	Size
OCT4 Concentrated	RM0148	1 ml
OCT4 Prediluted	RM0148RTU7	7 ml

IHC Procedure*

Positive Control Tissue: Seminoma
 Concentrated Dilution: 50-200
 Pretreatment: Tris EDTA pH9.0, 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human seminoma stained with anti-OCT4 using DAB

References:

1. Pseudogene OCT4-pg4 functions as a natural micro RNA sponge to regulate OCT4 expression by competing for miR-145 in hepatocellular carcinoma. Wang L, et al. Carcinogenesis 34:1773-81, 2013.
2. Identification of cancer stem-like side population cells in purified primary cultured human laryngeal squamous cell carcinoma epithelia. Wu CP, et al. PLoS One 8:e65750, 2013.
3. Breaking human cytomegalovirus major immediate-early gene silence by vasoactive intestinal peptide stimulation of the protein kinase A-CREB-TORC2 signaling cascade in human pluripotent embryonal Ntera2 cells. Yuan J, et al. J Virol 83:6391-403, 2009.