

Mouse Anti-ZAP70 [2F3.2]: MC0372

Intended Use: For Research Use Only

Description: ZAP-70 is a 70 kD protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. ZAP-70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP-70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP-70 negative) and Ig-unmutated (anti-ZAP-70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP-70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis.

Specifications

Clone: 2F3.2
Source: Mouse
Isotype: IgG2a
Reactivity: Human
Localization: Cytoplasm
Formulation: Tissue culture supernatant in PBS pH7.5, containing 0.2% BSA, 15mM sodium azide (NaN3)
Storage: Store at 2°- 8°C. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles
Applications: IHC
Package:

Description	Catalog No.	Size
ZAP70 Concentrated	MC0372	1 ml

IHC Procedure

Positive Control Tissue: Chronic lymphocytic
Concentrated Dilution: 25-200
Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
Incubation Time and Temp: 30-60 minutes @ RT
Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.

Doc. 100-MC0372
Rev. A