Medaysis

Rabbit Anti-p16/INK4a [CDKN2A/4844R]: RM0140, RM0140RTU7

Intended Use: For Research Use Only

Description: p16/INK4A is a tumor-suppressor protein and that genetic and epigenetic abnormalities in genes controlling the G1 checkpoint can lead to both escape from senescence and cancer formation. The interaction of p16/INK4 family members can be a binary complex with CDK4/6 or ternary complex with cyclin D-bound CDK4/6 and ultimately results in the inhibition of cell cycle progression. As such, expression of p16 INK4A is commonly associated with cellular senescence, and disruption of the p16 INK4A gene is frequently observed in human tumor. The p16/INK4A locus is deleted in a wide spectrum of tumors including melanoma, pancreatic adenocarcinoma, glioblastoma, certain leukemias, non-small cell lung cancer, cervical cancer, and bladder carcinoma.


Specifications

01	CDINIA / 40 4 4D		
Clone:	CDKN2A/4844R		
Source:	Rabbit		
Isotype:	IgG		
Reactivity:	Human		
Immunogen:	Purified recombinant p	rokaryotic full-length human	o16INK4a protein
Localization:	Nucleus and/or cytopla	sm	-
Formulation:	Antibody in PBS pH7.4	4, containing BSA and ≤ 0.099	% sodium azide (NaN3)
Storage:	Store at 2°- 8°C	-	
Applications:	IHC		
Package:			
Description		Catalog No.	Size
p16/INK4a Concentrated		RM0140	1 ml

p10/IINK4a Concentrated	KW0140	1 1111
p16/INK4a Prediluted	RM0140RTU7	7 ml

IHC Procedure*

Positive Control Tissue:	Squamous Cell Carcinoma	
Concentrated Dilution:	50-200	
Pretreatment:	Tris EDTA pH9.0, 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C	
Incubation Time and Temp:	30-60 minutes @ RT	
Detection:	Refer to the detection system manual	
* Result should be confirmed by an established diagnostic procedure.		



FFPE human cervix stained with anti-p16 using DAB

References:

- 1. Inhibition of the 60S ribosome biogenesis GTPase LSG1 causes endoplasmic reticular disruption and cellular senescence. Pantazi A, et al. Aging Cell. Aug;18(4):e12981, 2019.
- 2. CDKN2A copy number and p16 expression in malignant pleural mesothelioma in relation to asbestos exposure. Eeva Kettunen, et al. BMC Cancer volume 19, Article number: 507, 2019.
- 3. Protein and chemotherapy profiling of extracellular vesicles harvested from therapeutic induced senescent triple negative breast cancer cells. Kavanagh EL, et al. Oncogenesis. Oct 9;6(10):e388, 2017.

Doc. 100-RM0140 Rev. A