Enable Innovation DATA SHEET

Rabbit Anti-MAP3K3/MEKK3 (Mitogen-Activated Protein Kinase Kinase Kinase 3) Polyclonal: RC0318

Intended Use: For Research Use Only

Description: MEKKs (Mitogen activated protein kinase kinases) are serine-threonine kinases that act as the first tier of cellular MAP kinase pathways by activation of MAP/ERK kinases, or MEKs. Many enzymes with MEKK activity have been identified, including MEKK1-4, Raf, MLK3, TAK, and DLK. MEKKs generally display little similarity outside of their catalytic kinase domains. MEKK1-4 are nearly 50% identical within their catalytic domains, and are known to regulate Erk, Jnk, and p38 MAP kinase pathways. MEKK2 and MEKK3 bind MEK5 via conserved PB1 domains, leading to downstream activation of Erk5.

Specifications

Clone: Polyclonal Source: Rabbit Isotype: IgG

Reactivity: Human, mouse

Immunogen: Synthesized peptide derived from the internal region of human MEK Kinase-3

Localization: Cytoplasm

Formulation: Antibody in PBS pH7.4, containing BSA and ≤ 0.09% sodium azide (NaN3)

Storage: Store at 2°-8°C

Applications: IHC, ELISA, ICC/IF, WB

Package:

Description	Catalog No.	Size	
MAP3K3/MEKK3 (Mitogen-Activated Protein Kinase Kinase	RC0318	1 ml	
Kinase 3) Concentrated			

IHC Procedure


Positive Control Tissue: Ovary carcinoma, HeLa cells

Concentrated Dilution: 10-50

Pretreatment: Tris EDTA pH9.0, 15 minutes ressure Cooker or 30-60 minutes water bath at 95°-99°C

Incubation Time and Temp: Overnight @ 4°C

Detection: Refer to the detection system manual * Result should be confirmed by an established diagnostic procedure.



FFPE human lung cancer stained with anti-MEKK3 using DAB

References:

- 1. PLX4032, a selective BRAF(V600E) kinase inhibitor, activates the ERK pathway and enhances cell migration and proliferation of BRAF melanoma cells. Halaban R, et al. Pigment Cell Melanoma Res 23:190-200, 2010.
- 2. Phosphorylation of MEKK3 at threonine 294 promotes 14-3-3 association to inhibit nuclear factor kappaB activation. Matitau AE & Scheid MP. J Biol Chem 283:13261-8, 2008.
- 3. Expression profiles of protein tyrosine kinase genes in human embryonic stem cells. Son MY, et al. Reproduction 136:423-32, 2008.

Doc. 100-RC0318

Rev. A

Orders: customercare@medaysis.com Support: techsupport@medaysis.com Tel: 510-509-3153 www.medaysis.com www.medaysis.com</