

Mouse Anti-FOXL1 [PCRP-FOXL1-1F8]: MC0474, MC0474RTU7

Intended Use: For Research Use Only

Description: Forkhead box protein L1s or FKHL11, is encoded by the FOXL1 gene in human. FOXL1 protein belongs to the forkhead box (Fox) family of transcription factors characterized by a common DNA-binding domain (the forkhead box). FOXL1 is involved in regulating the proliferation of gastrointestinal epithelial cells. Loss of Foxl1 leads to distorted architecture of the stomach and small intestine in mice due to a markedly increased epithelial proliferation. Foxl1 knockout is correlated with an enhanced Wnt/ β -catenin pathway activation as evidenced by increased β -catenin nuclear localization. FOXL1 expression is down-regulated in the majority of human clear cell renal cell carcinoma (ccRCC), at both mRNA and protein levels. FOXL1 expression inversely correlates with the aggressive phenotype of ccRCC and the survival outcome of patients.

Specifications


Clone: PCRP-FOXL1-1F8
 Source: Mouse
 Isotype: IgG1
 Reactivity: Human
 Immunogen: Recombinant fragment aa 104-258 of human FOXL1 protein
 Localization: Nucleus
 Formulation: Purified antibody in PBS pH7.4, containing BSA and \leq 0.09% sodium azide (NaN₃)
 Storage: Store at 2°- 8°C
 Applications: IHC, Flow Cyt., ICC/IF, IP, WB
 Package:

Description	Catalog No.	Size
FOXL1 Concentrated	MC0474	1 ml
FOXL1 Prediluted	MC0474RTU7	7 ml

IHC Procedure*

Positive Control Tissue: Testis, muscle, breast cancer
 Concentrated Dilution: 50-200
 Pretreatment: Tris EDTA pH9.0, 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human prostate stained with anti-FOXL1 using DAB

References:

1. Low level of FOXL1 indicates a worse prognosis for gastric cancer patients. Ertao Z, et al., Tumour Biol. Aug; 37(8): 11331-7, 2016.
2. Inhibitory effects of forkhead box L1 gene on osteosarcoma growth through the induction of cell cycle arrest and apoptosis. Chen X, et al. Oncol Rep. Jul;34(1):265-71, 2015.
3. Forkhead box L1 is frequently downregulated in gallbladder cancer and inhibits cell growth through apoptosis induction by mitochondrial dysfunction. Qin Y, et al, PLoS One. Jul 10;9(7):e102084, 2014.