

Rabbit Anti-Paxillin [MD380R]: RM0159

Intended Use: For Research Use Only

Description: Paxillin is a cytoskeletal protein involved in actin-membrane attachment at sites of cell adhesion to the extracellular matrix (focal adhesion). It is a multidomain protein. The C-terminal region of paxillin contains four LIM domains that target paxillin to focal adhesions, presumably through a direct association with the cytoplasmic tail of beta-integrin. The N-terminus of paxillin controls most of its signaling activity. The proteins that bind to paxillin are diverse and include protein tyrosine kinases, such as Src and FAK, structural proteins, such as vinculin and actopaxin, and regulators of actin organization, such as COOL/PIX and PKL/GIT. Paxillin is widely expressed in epithelial cells of various tissues, neuronal cells and mesenchymal derived cells. An antibody to Paxillin is helpful in differentiating between renal cell carcinoma (Paxillin negative) and chromophobe renal cell carcinoma or renal oncocytoma (Paxillin positive), which are rare renal tumors originating from the intercalated cells of collecting ducts. Paxillin has been reported to be involved in tumor invasion and metastasis. Its expression in lung and liver cancers has been correlated with advanced tumor stage and metastasis.


Specifications

Clone:	MD380R
Source:	Rabbit
Isotype:	IgG
Reactivity:	Human
Immunogen:	Peptide corresponding to the N-terminus of human Paxillin
Localization:	Cytoplasm
Formulation:	Antibody in PBS pH7.4, containing BSA and $\leq 0.09\%$ sodium azide (NaN ₃)
Storage:	Store at 2°- 8°C
Applications:	IHC, IF, WB
Package:	

Description	Catalog No.	Size
Paxillin Concentrated	RM0159	1 ml

IHC Procedure*:

Positive Control Tissue:	Breast cancer, tonsil
Concentrated Dilution:	25-100
Pretreatment:	Tris EDTA pH9.0, 15 minutes Pressure Cooker or 30-60 minutes water bath at 95°-99°C
Incubation Time and Temp:	30-60 minutes @ RT
Detection:	Refer to the detection system manual
* Result should be confirmed by an established diagnostic procedure.	



FFPE human tonsil stained with anti-Paxillin using DAB

References:

1. Immunohistochemical expression of paxillin in potentially malignant disorders and squamous cell carcinoma patients
2. Shakir Alam, et al. J Oral Maxillofac Pathol. Oct 17;26(3):322-329, 2022.
3. Paxillin: a crossroad in pathological cell migration. Ana María López-Colomé, et al. Journal of Hematology & Oncology volume 10, Article number: 50, 2017.
4. A paxillin tyrosine phosphorylation switch regulates the assembly and form of cell-matrix adhesions. Ronen Zaidel-Bar, et al. J Cell Sci. 120 (1): 137-148, 2007.

Doc. 100-RM0159
Rev. A